

Spin-Charge Separation and Kinetic Energy in the t - J Model

Sanjoy K. Sarker

Department of Physics and Astronomy
The University of Alabama, Tuscaloosa, AL 35487

February 2, 2008

Abstract

The effect of spin-charge separation on the kinetic energy of the two-dimensional t - J model is examined. Using a sum rule, we derive an exact expression for the lowest possible KE (E_{bound}) for any state without doubly occupied sites. The kinetic energies of the relevant slave-boson and Schwinger-boson mean-field states are found to be considerably larger than E_{bound} . The MF states exhibit complete spin-charge separation, and form the basis of a number of microscopic theories. An examination of the momentum distribution reveals that the large increase in KE of the MF states is due to excessive depletion of electrons from the bottom of the band (Schwinger boson states) and holes from the top of the band (slave boson states). To see whether the excess KE is simply due to the poor treatment of the local constraints, we solve the constraint problem analytically for the Schwinger boson MF states in the $J = 0$ limit. This restores gauge invariance, incorrectly violated in the MF theories. The resulting state is a generalization of the Hartree-Fock state in the Hubbard model, but one that includes spin-wave excitations, removing a deficiency of the simple HF theory. Even after the constraints are imposed correctly, the MF kinetic energy is found to be much larger than E_{bound} . These results support the notion, advanced in earlier papers, that spin-charge separation in the MF state costs excessive kinetic energy, and makes the state unstable toward recombination processes which lead to superconductivity in $d = 2$ and a Fermi liquid state in higher dimensions.

PACS: 71.10 Fd, 74.25 -q, 74.72 -h, 74.62 Dh

I. INTRODUCTION

The possibility of spin-charge separation in large- U Hubbard model and the equivalent t - J model is of considerable interest because of its relevance to high- T_c superconductivity [1]. Such a behavior is seen in $d = 1$, where the metallic state is characterized by separate spin and charge excitations, not electron-like quasiparticles [2]. The metallic state evolves out of the insulating state at half filling, which has no magnetic order. By contrast, for $d \geq 2$, the ground state is antiferromagnetically ordered at half filling. Therefore, if the 2-d Hubbard model were to describe the cuprates, its low-energy behavior would be governed by several distinct fixed points: an AF insulator; a metallic, presumably spin-charge separated state without magnetic order; possibly a superconducting state; and possibly even a Fermi liquid at lower electron densities.

A number of microscopic theories are based on the slave-boson (SLB) [3, 4] and the Schwinger-boson (SWB)[5] representations of the 2-d t - J model in which the electron field is decomposed into a spin (spinon) and a charge (holon) component, subject to a local constraint. Each theory is centered on a mean-field state which is supposed to take care of short-range or high-energy processes, leaving behind weak residual interactions. The MF state is characterized by independent spinon and holon excitations, and thus exhibits complete spin-charge separation. The results have been varied, but not too satisfactory. That none of the MF theories succeed suggests that there is a generic reason for the failure – some important bit of physics is missing in all of them. Identifying it and understanding the energetics behind the failure is of great interest.

One obvious source of error, to be considered later, is the local constraint, which is treated on the average in all MF theories. We have argued previously on continuity grounds that there is another, and more fundamental, reason for the failure [6, 7]. The MF states exhibit maximum spin-charge separation

in all dimensions and at all densities. This can not be true, since in high dimensions (e.g., in $d = 3$), one expects a Fermi liquid state to emerge. In $d = 1$, strong spin-charge separation occurs because electrons of opposite spins can not pass each other, and thus are localized as “spins” or “local moments”. This costs extra kinetic energy. In higher dimensions, electrons can avoid double occupancy by going around each other and delocalize, and thereby reduce the kinetic energy. Therefore, spin-charge separation should weaken with increasing dimensionality and decreasing electron density. MF states fail because they show too much spin-charge separation which costs excessive kinetic energy. They would be unstable toward spin-charge recombination into physical electrons, the latter would try to form a Fermi liquid. Recombination will help destroy magnetic order because, as the electrons delocalize, the size of the moment is reduced. It will be more effective at higher dimensions and far from half filling since more pathways are available, which would explain the emergence of the Fermi liquid state in $d = 3$.

Spin-charge recombination is a collective process which appears through the kinetic energy term. It has been analyzed in $d = 2$ using RPA in the Schwinger-boson representation. The electron is a “collective” excitation. Recombination is strong enough to destroy magnetic order, but the normal state remains weakly spin-charge separated [6]. Additional spin-charge recombination, in which a pair of holons combine with a pre-paired spin singlets then leads to d -wave superconductivity [7]. In this picture the destruction of magnetic LRO, the appearance of superconductivity and the emergence of a Fermi liquid state at high dimensions are all caused by spin-charge recombination, as the system evolves away from half filling.

In this paper we present two exact results which allows us to get more insight into the energetics of spin-charge separation. Specifically, we examine the hypothesis that MF states have excessive kinetic energy. This issue is quite subtle. How does one know how much KE is too much? With increasing U , kinetic energy of the actual ground state will increase, as electrons are excited out of the Fermi sea. As $U \rightarrow \infty$, there is no double occupancy, and we expect KE to be quite large (compared with the noninteracting case, E_{free}), even if the state is a Fermi liquid. By using a sum rule we determine the lowest possible kinetic energy (E_{bound}) for any state of the t - J model, i.e., for any state in which there is no double occupancy. As expected, we find that $E_{bound} \gg E_{free}$. Interestingly, near half filling, the Schwinger-boson state has a lower KE than the slave-boson state. However, as we shall see, the kinetic energy of either state is much larger than E_{bound} . An examination of the momentum distribution reveals that excess KE is due to excessive depletion of single particle states near the band edges – a situation that is quite likely to give rise to strong fluctuations.

The second issue involves the largely unsolved problem of the constraints. Treating them on the average would be valid if fluctuations were small enough to be treated perturbatively. However, a perturbation theory is not likely to converge, and may make matters worse. The constraints imply a local gauge symmetry which, according to Elitzur’s theorem [8], can not be spontaneously broken. MF theories break this symmetry so that elementary excitations, i.e., spinons and holons, are not gauge invariant and therefore not observable. The true ground state and elementary excitations are of course gauge invariant. Since gauge invariance can not be restored perturbatively it is hard to judge the correctness of the theory, even at a qualitative level.

One way to avoid the constraint problem is to consider the Hartree-Fock (HF) states of the Hubbard model [9], in which constraints are treated exactly. These states are related to the Schwinger boson MF states. But the HF theory has its own shortcomings. The spin-wave excitations are missing at the HF level, so that low-energy behavior is not described correctly. For example, magnetic order persists all the way upto a $T_c \sim U$ in all dimension, violating the Marmin-Wagner theorem. On the other hand, for the Heisenberg model, the Schwinger boson MF theory is qualitatively similar to the spin-wave theory, which is based on the Holstein-Primakoff (HP) representation and is gauge invariant [10, 11]. But HP theory applies only to half-filling, there is no charge excitations.

Here we solve these problems in the $U = \infty$ limit. This limit is of interest because the mapping onto the t - J model is exact, so a direct comparison between SWB and HF results can be made. By integrating out the constraints in the Schwinger boson theory, we obtain a generalized HP representation which is valid away from half filling. This, in turn, leads to a generalized Hartree-Fock Hamiltonian, which include both charge and spin excitations. The procedure restores gauge invariance, and the inclusion of the spin-waves takes care of the deficiencies of the HF theory. We find that even after the constraints are taken into account, the nature of the state is not changed, nor is the degree of spin-charge separation. Kinetic energy is reduced,

but continues to be excessive compared with E_{bound} , which supports our hypothesis.

II. THE MODEL AND SPIN-CHARGE SEPARATION

We start with a discussion of the meaning of spin-charge separation. The Hubbard model is characterized by nearest-neighbor hopping t , and on-site repulsion U . For large U , doubly occupied sites can be projected out, which leads to the t - J Hamiltonian:

$$H = -t \sum_{ij} (c_{i\sigma}^\dagger c_{j\sigma} + h.c.) + J \sum_{ij} (\mathbf{S}_i \cdot \mathbf{S}_j - \frac{1}{4} n_i n_j), \quad (1)$$

where the summation is over nearest neighbors. The second term describes superexchange, with $J = 4t^2/U$. The first term describes hopping, and has the same form as that in the Hubbard model. However, $c_{i\sigma}^\dagger$ creates a “projected electron”, i.e., an electron of spin σ , only if the site i is unoccupied. At half-filling, one needs to consider only the exchange term.

In $d = 1$, the Hubbard model has been diagonalized by Lieb and Wu [12], using the Bethe Ansatz. The occurrence of spin-charge separation can be seen most clearly in the large U limit where, as shown by Ogata and Shiba [13], the ground-state wave function for N_e electrons can be approximately written as:

$$\psi = \psi_{charge} \psi_{spin}. \quad (2)$$

Here $\psi_{charge}(x_1, x_2, \dots)$ is a determinantal wave function for N_e *spinless* fermions, where x_i are the electron coordinates, irrespective of spin. And ψ_{spin} is the Bethe’s solution for the Heisenberg model for N_e spins (with the holes removed). The wave function at smaller U is more complicated, but is continuously connected to the large U solution in the renormalization group sense. Of course, the low-lying excited states also similarly decompose – i.e., the system is characterized by independent spin and charge excitations. A good account of the meaning of Eq. (2) and its consequences is given by Anderson [2].

Usually spin-charge separation is associated with the Luttinger liquid state, which is a metal and has no magnetic order. However, Eq. (2) clearly suggests that the phenomenon is more general. Any wave function which has the form of (2), or is continuously connected to such a wave functions, exhibits spin-charge separation. Consider the following cases.

a) Half-filled case: If we continue the Lieb-Wu wave functions to half-filling, the form (2) will clearly be preserved. The (spinless) charge band becomes filled, and the system turns into an insulator. A moment’s reflection shows that such spin-charge separation occurs at half filling in all dimensions. The reduction of the large- U Hubbard model to the Heisenberg model is precisely due to spin-charge decoupling. There is exactly one charge per site, which can be viewed as a filled band of spinless fermions. For $d \geq 2$, the insulating state is magnetically ordered.

b) Mean-Field States: Similarly, mean-field wave functions for the t - J model also exhibits spin-charge separation. In this case the electron operator is represented as:

$$c_{i\sigma}^\dagger = b_{i\sigma}^\dagger h_i,$$

where $b_{i\sigma}^\dagger$ creates a spinon and h_i creates a holon, subject to the constraint

$$b_{i\uparrow}^\dagger b_{i\uparrow} + b_{i\downarrow}^\dagger b_{i\downarrow} + h_i^\dagger h_i = 1.$$

The spin-operators are then represented as: $S_i^+ = b_{i\uparrow}^\dagger b_{i\downarrow}$, and $S_i^z = \frac{1}{2}(b_{i\uparrow}^\dagger b_{i\uparrow} - b_{i\downarrow}^\dagger b_{i\downarrow})$. In the Schwinger boson representation, the spinon is a boson and holon, a fermion. The opposite is true in the slave boson representation. In either case, constraints are treated on the average, and a MF approximation leads to the Hamiltonian:

$$H_{MF} = H_{spinon} + H_{holon}. \quad (3)$$

Note that all eigenstates of H_{MF} are of the form of Eq.(2). This is also true in the presence of arbitrary spinon-spinon and holon-holon interactions.

In general bosons condense, leading to long-range order (LRO). In the Schwinger-boson case, the ordering is magnetic. The mean-field theory gives quite a good description of the Neel state at half filling [10]. Away from half filling, the AF insulator evolves continuously into a metallic state with a spiral LRO, and eventually to a ferromagnetic metal at small J/t [5]. In the metallic region, ψ_{charge} in the SWB theory is again a spinless fermion determinant.

c) Hartree-Fock states: The magnetically ordered states obtained by the Hartree-Fock treatment of the Hubbard model [9] constitute a third example of spin-charge separation. The HF decomposition of the U term leads to the quadratic Hamiltonian

$$H_{HF} = H_0 - 2U \sum_i \mathbf{m}_i \cdot \mathbf{S}_i, \quad (4)$$

where H_0 is the hopping part and $\mathbf{m}_i = \langle \mathbf{S}_i \rangle$ is the local magnetization, which is determined self consistently. To describe spiral states, it is convenient to break the spin rotational symmetry in the xy plane and choose $\langle S_i^+ \rangle = m e^{iQ \cdot r}$ as the order parameter, where Q is the spiral wavevector which continuously connects the AF state ($Q = (\pi, \pi)$) at half-filling, to the ferromagnetic state ($Q = 0$) which occurs at large $U\delta/t$. The theory has been described in detail elsewhere [9]. The HF approximation leads to two magnetic bands separated by a gap of size $2Um$. At large U , low-energy states are confined to the lower band. However, there is no spin-degeneracy; each one-particle state can accomodate one electron, not two. Therefore, low-energy wave functions are one-component (i.e., spinless) determinantal wave functions, similar to ψ_{charge} in Eq. (2). The HF states are spin-charge separated because they are closely related to the corresponding states of the Schwinger boson MF theory.

To see this, consider H_{HF} for $Um \gg t$. Then, the spin of an electron will locally be parallel to the local moment $\mathbf{m}_i$. Let $f_i^\dagger$ and $g_i^\dagger$ create electrons with spin parallel and antiparallel to $\mathbf{m}_i$, respectively. The bare electron operators, which are quantized along the z direction, can be expanded as:

$$a_{i\sigma} = u_{i\sigma} f_{i\sigma} + \text{sgn}(\sigma) u_{i,-\sigma}^* g_{i\sigma}, \quad (5)$$

where $\sigma = 1$ for $\uparrow$ and -1 for $\downarrow$. The complex numbers $u_{i\sigma}$ depend on the direction of $\mathbf{m}_i$, and satisfy the normalization condition

$$\sum_{\sigma} u_{i\sigma}^* u_{i\sigma} = 1, \quad (6)$$

Suppose, $\mathbf{m}_i = (m, \theta, \phi)$, then a simple representation is:

$$u_{i\uparrow} = \cos(\theta_i/2), \quad u_{i\downarrow} = \sin(\theta_i/2) e^{i\phi_i}.$$

Note that the second term in H_{HF} is diagonal in the f, g representation:

$$-Um \sum_i (f_i^\dagger f_i - g_i^\dagger g_i).$$

For $U \rightarrow \infty$, we can neglect the g terms which constitute the upper band. The similarity with the Schwinger boson representation is obvious, with $u_{i\sigma}$ playing the role of $b_{i\sigma}$ and f_i the charge part of the electron operator. The holon operator is obtained by a particle-hole transformation: $f_i \rightarrow h_i^\dagger$. Eq. (6) now plays the role of the constraints. The difference is that b 's are operators, where as u 's are c-numbers. However, in the MF approximation, b 's condense, and we can take $u_{i\sigma} \approx \langle b_{i\sigma} \rangle$. For $U = \infty$, the ground state is a ferromagnetic metal away from half filling. Then $\mathbf{m}_i = \mathbf{m}$, (and hence $u_{i\sigma} = u_\sigma$), is independent of i . The HF wave function, written in terms of the original electron coordinates, (those created by bare operators $a_{i\sigma}$), spin-charge separates, with ψ_{charge} again given by the one component Slater determinant, and is independent of $\mathbf{m}$. ψ_{spin} describes the condensed part and is just a constant.

For finite but large U , there will be small admixture of g 's via the hopping term which can be eliminated perturbatively, to obtain contribution of order J — the exchange contribution. However, as in the t - J model,

such corrections are due to spin-spin interaction and thus does not change our conclusion about spin-charge separation. There are of course some important differences between the two theories. In the SWB MF theory, constraints are treated on the average, whereas in HF theory they are treated exactly. Similarly, in the HF theory, the low energy spin excitations are absent, and hence physics of the singlets is not included.

Spin-Charge Separation and Local Moments

At half-filling, it is natural to associate spin-charge separation with the formation of local moments. In $d = 1$, evidently the local moment picture persists away from half-filling since ψ_{spin} continues to be the wave function of a “squeezed” Heisenberg antiferromagnet [13, 2]. At large U , two electrons of opposite spin can not go past each other. Hence, electrons are localized within a region of size $(1 - \delta)^{-1}$, and the size of the moment is approximately $(1 - \delta)/2$. On the other hand, if they can delocalize there will be no moment, nor will there be spin-charge separation. Therefore we can associate spin-charge separation with “localized” moments. Evidently this picture is consistent with Schwinger boson MF states, as well as HF states. Note that spin-charge separation is associated with the amplitude, and does not require the existence of LRO (phase coherence). The latter disappears at any finite T in $d = 2$ (Mermin-Wagner theorem). But the wave-function remains spin-charge separated since, at the MF level, the moment vanishes only at $T \sim U$. Because moment formation involves a type of localization, it would cost kinetic energy. Interestingly, from this point of view a transition from a Luttinger liquid to a Fermi liquid can be considered a form of delocalization transition.

III. SPECTRAL SUM-RULE AND LOWER BOUND ON KINETIC ENERGY

In this section we obtain an exact expression for the lowest possible KE in the t - J model. It is useful to express the hopping part of the t - J Hamiltonian in the momentum space

$$H_{hop} = \sum_{\mathbf{k}, \sigma} \epsilon(\mathbf{k}) c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma}, \quad (7)$$

where $\epsilon(\mathbf{k}) = -2t(\cos k_x + \cos k_y)$ is the free-electron hopping energy, and $c_{\mathbf{k}\sigma}^\dagger$ is the projected electron operator in the momentum space. The average hopping energy (per site) is then given by

$$E_{kin} = \frac{1}{N} \sum_{\mathbf{k}} \epsilon(\mathbf{k}) n_c(\mathbf{k}), \quad (8)$$

where $n_c(\mathbf{k}) = \sum_{\sigma} n_{c\sigma}(\mathbf{k})$ is the (total) electron momentum distribution, with $n_{c\sigma}(\mathbf{k}) = \langle c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} \rangle$. Eq.(8) is exact. Hence any errors in E_{kin} are due to approximations used in calculating $n_c(\mathbf{k})$. The kinetic energy of the Hubbard model is given by the same formula, except that the corresponding momentum distribution differs from $n_c(\mathbf{k})$ by terms of order J , and $J\delta$. Here, we focus on the kinetic energy as defined above. Of course, for $U = \infty$ (i.e., $J = 0$), the mapping is exact.

Usually one is interested in the singularities of $n_c(\mathbf{k})$ near the Fermi surface. However, in the present case, we need to know $n_c(\mathbf{k})$ for all $\mathbf{k}$. In order to minimize the kinetic energy, the system will try to make $n_c(\mathbf{k})$ as large as possible near the bottom of the noninteracting band $\epsilon(\mathbf{k})$, and as small as possible near the top. Hence states with $\mathbf{k}$ near the band edges are quite important. This has serious implications for the mean-field states. The assumption underlying the microscopic theories is that high energy processes, i.e., those involving electrons with $\mathbf{k}$ far from the Fermi sea, are accurately taken care of at the mean-field level, leaving behind residual interactions which describes low-energy physics. Therefore, if a MF state does not yield accurate values of $n_c(\mathbf{k})$ near the band edges, it is likely to be unstable toward fluctuations which transfer electrons from the top to the bottom of the band, and thereby lower the kinetic energy considerably.

A. Spectral Sum Rule:

Now, the noninteracting state has the the lowest KE, with $n_c(\mathbf{k}) = 2$ below the Fermi level ϵ_F . However, this is not a good reference state since any interaction would increase the kinetic energy by exciting electrons

out of the Fermi sea and, as we see below, at large U , a substantial part of the increase comes from avoiding double occupancy, which may have nothing to do with spin-charge separation. Fortunately, we can determine the lowest possible kinetic energy in the t - J model. Since the latter does not have double occupancy, the corresponding increase in KE is accounted for at the outset.

Formally, we have

$$n_{c\sigma}(\mathbf{k}) = \int \frac{dz}{2\pi} A_{c\sigma}(\mathbf{k}, z) f(z), \quad (9)$$

where $A_{c\sigma}(\mathbf{k}, \omega)$ is the spectral function for projected electrons which obeys a sum rule [14]. Here we give a derivation which is useful for this paper. Let us define the momentum distribution for physical holes (not to be confused with holons): $p_{c\sigma}(\mathbf{k}) \equiv \langle c_{\mathbf{k}\sigma} c_{\mathbf{k}\sigma}^\dagger \rangle$. This is obtained from

$$p_{c\sigma}(\mathbf{k}) = \int \frac{dz}{2\pi} A_{c\sigma}(\mathbf{k}, z) (1 - f(z)). \quad (10)$$

Hence, total spectral weight for spin σ is given by

$$\int \frac{dz}{2\pi} A_{c\sigma}(\mathbf{k}, z) = n_{c\sigma}(\mathbf{k}) + p_{c\sigma}(\mathbf{k}) = \langle \{c_{\mathbf{k}\sigma}, c_{\mathbf{k}\sigma}^\dagger\} \rangle. \quad (11)$$

For usual fermions operators, the anticommutator equals unity, giving the usual spectral sum rule. This is not true for projected electrons. To evaluate the right hand side, we use the representation of the electron operator in terms of spinon and holon operators which, in momentum space, gives

$$c_{\mathbf{k}\sigma}^\dagger = N^{-\frac{1}{2}} \sum_{\mathbf{q}} b_{\mathbf{k}+\mathbf{q},\sigma}^\dagger h_{\mathbf{q}}.$$

Then the average value of the anticommutator is given by

$$\langle \{c_{\mathbf{k}\sigma}, c_{\mathbf{k}\sigma}^\dagger\} \rangle = \frac{1}{N} \sum_i (\langle b_{i\sigma}^\dagger b_{i\sigma} \rangle + \langle h_i^\dagger h_i \rangle) = n_{b\sigma} + \delta, \quad (12)$$

where $n_{b\sigma}$ is the average number spinons (per site) of spin σ , and δ is the average number holons per site. It is useful to separate into a charge and a spin part. Let $n_{b\sigma} = n_b/2 + \sigma m$, where $m = (n_{b\uparrow} - n_{b\downarrow})/2$ is the average magnetization, and $n_b = \sum_{\sigma} n_{b\sigma} = 1 - \delta$ (from the constraint condition). Then we have

$$n_{c\sigma}(\mathbf{k}) + p_{c\sigma}(\mathbf{k}) = \frac{1}{2}(1 + \delta) + \sigma m. \quad (13)$$

We stress that these sum rules are exact. Since the right hand side depends on the spinon and holon densities the sum rules are valid in either representation, and in all dimensions, irrespective of whether the local constraint is imposed exactly or on the average. Summing over σ we obtain the total electron spectral weight:

$$n_c(\mathbf{k}) + p_c(\mathbf{k}) = (1 + \delta). \quad (14)$$

The remaining $(1 - \delta)$ resides in the upper Hubbard band. Note that, in contrast to bare electrons, the spectral weight (Eq. 13) depends on charge and spin densities.

The sum rule, Eq. (14), implies that each momentum state $\mathbf{k}$ can accomodate at most $1 + \delta$ electrons (compared with 2 for a noninteracting system). It follows that, at half filling, in order to accomodate all the electrons, one must have $n_c(\mathbf{k}) = 1$ for all $\mathbf{k}$, which is the correct result. In the Schwinger boson MF theory this can be viewed as a filled holon band. The kinetic energy per electron is maximum ($= 0$), as compared with $\sim -t$, for free electrons.

B. Lower Bound for the Kinetic Energy:

Away from half filling, the exact $n_c(\mathbf{k})$ is not known. But it is clear that states with low kinetic energy will have $n_c(\mathbf{k})$ large near the bottom of the band and small near the top. Indeed, the lowest possible kinetic

energy for the model is obtained by filling up a *modified Fermi sea* with exactly $n_c(\mathbf{k}) = (1 + \delta)$, upto a new Fermi level ϵ_{F1} . Above ϵ_{F1} , $n_c(\mathbf{k}) = 0$. This gives an exact lower bound for the kinetic energy per site.

$$E_{bound} = \frac{1 + \delta}{N} \sum_{\mathbf{k}} \epsilon(\mathbf{k}) \theta(\epsilon_{F1} - \epsilon(\mathbf{k})), \quad (15)$$

where ϵ_{F1} is determined from

$$n_c = \frac{1 + \delta}{N} \sum_{\mathbf{k}} \theta(\epsilon_{F1} - \epsilon(\mathbf{k})).$$

where $n_c = 1 - \delta$. The modified Fermi sea is obtained from the noninteracting one by exciting $1 - \delta$ electrons from each state below the (noninteracting) Fermi level ϵ_F to a state above. Therefore $\epsilon_{F1} > \epsilon_F$, and the kinetic energy is correspondingly larger (see Fig. 1). This cost in kinetic energy is due to the removal double occupancy. In other words, E_{bound} is the lowest possible kinetic energy for any state in which double occupancy is forbidden.

Apart from the prefactors, the expression for E_{bound} at a density $n_c = 1 - \delta$ is identical to that of a noninteracting system at a higher density $n' = 2n_c/(1 + \delta)$:

$$E_{bound}(n_c) = \frac{1 + \delta}{2} E_{free}\left(\frac{2n_c}{1 + \delta}\right). \quad (16)$$

The volume of the modified Fermi sea is $2/(1 + \delta)$ times the free electron Fermi sea.

What sort of a state corresponds to E_{bound} ? Since it evidently resembles a free-fermion state with a sharp Fermi surface, let us consider the following wave function:

$$|\Psi\rangle = \mathcal{Z} \prod_{k < k_F, \sigma} c_{\mathbf{k}\sigma}^\dagger |vac\rangle,$$

where $\mathcal{Z}$ is a normalization constant. Since $c_{\mathbf{k}\sigma}$ represents a projected electron, it is easy to see this is the Gutzwiller wavefunction in the infinite- U limit. Now, the projected electron operators for arbitrary spins satisfy the following commutation rules

$$\{c_{\mathbf{k}\sigma}, c_{\mathbf{k}',\sigma'}^\dagger\} = \frac{1}{N} \sum_i e^{i(\mathbf{k}' - \mathbf{k}) \cdot \mathbf{r}_i} [b_{i\sigma'}^\dagger b_{i\sigma} + \delta_{\sigma\sigma'} h_i^\dagger h_i],$$

Notice that right-hand side, which arises from the on-site anticommutators, consists of Fourier components of charge and spin-density operators. Suppose, as an approximation, we replace the right-hand side by its average value and assume that there is no magnetic or CDW order. Then, we obtain

$$\{c_{\mathbf{k}\sigma}, c_{\mathbf{k}',\sigma'}^\dagger\} \approx \xi \delta_{\sigma,\sigma'} \delta_{\mathbf{k}\mathbf{k}'},$$

where $\xi = (1 + \delta)/2$. In other words, $c_{\mathbf{k}\sigma}$ is effectively replaced by $\xi^{\frac{1}{2}}$ times an usual fermion operator which carries spin. If k_F is chosen to yield the correct electron density in the same approximation, then the new Fermi surface agrees with the modified one, and the energy is given by E_{bound} . Since spin- and charge-density fluctuations are completely suppressed, this state does not have any localized moment, and hence, there is no spin-charge separation.

IV. KINETIC ENERGY OF THE MEAN-FIELD STATES

The kinetic energy of an actual state is greater than E_{bound} . Now, the momentum distribution for the MF states is given by a convolution:

$$n_c(\mathbf{k}) = \frac{1}{N} \sum_{\mathbf{q}} n_b(\mathbf{k} + \mathbf{q})(1 \pm n_h(\mathbf{q})), \quad (17)$$

where $n_b(\mathbf{k})$ and $n_h(\mathbf{k})$ are the momentum distribution functions for spinons and holons, respectively. The minus (plus) sign corresponds to the Schwinger (slave) boson representation. Consider the MF states at $T = 0$.

a) *Slave-boson State*: We consider an RVB type state in which bosonic holons condense at $\mathbf{k} = 0$, giving

$$n_c(\mathbf{k}) = 1 - \delta(1 - n_b(\mathbf{k})). \quad (18)$$

Spinons form their own Fermi sea, identical to that for noninteracting electrons, and $n_b(\mathbf{k}) = 2$ for occupied and 0 for unoccupied states. Hence, electrons have the same Fermi surface, but $n_c(\mathbf{k})$ equals $1 + \delta$ below, and $1 - \delta$ for all $\mathbf{k}$ above the Fermi level all the way to the top of the band. Since $\sum_{\mathbf{k}} \epsilon(\mathbf{k}) = 0$, the kinetic energy per site is given by

$$E_{slave} = \delta E_{free}(1 - \delta). \quad (19)$$

b) *Schwinger-boson State*: We consider the ferromagnetic state (small J) since the kinetic energy is lowest in this case. (At larger J , the kinetic energy, as defined here, is larger due to the competition with antiferromagnetic fluctuations). Spinons condense at $\mathbf{k} = 0$, so that

$$n_c(\mathbf{k}) = (1 - \delta)(1 - n_h(-\mathbf{k})). \quad (20)$$

The holons form a Fermi surface, centered at the zone corner, with $n_h(\mathbf{k}) = 1$ in the occupied region, and zero in the unoccupied region. The electrons have the same Fermi surface, so the electron Fermi energy is much higher than the noninteracting one. On the average, there are only $1 - \delta$ electrons below this Fermi level (even less than the number at half filling!) and none above. In this case the kinetic energy per site is

$$E_{sch} = \frac{(1 - \delta)}{2} E_{free}(2\delta). \quad (21)$$

Fig (1) and (2) shows $n_c(\mathbf{k})$ and the kinetic energy for various states. First, note that all states have the correct kinetic energy (= zero) at half filling. Away from half filling, KE for either MF state is much larger than E_{bound} , by about a factor of two in the small- δ region. Surprisingly, for $\delta < 1/3$, the Schwinger boson state has a lower kinetic energy than the slave boson state. One expects Schwinger boson theory to do better as far as exchange energy is concerned since it works well for the Heisenberg model. It appears that it is also better for the KE at low doping, the region of interest for the cuprates. In the slave boson case, the Fermi surface coincides with the noninteracting one, and below the Fermi level, $n_c(\mathbf{k})$ is the maximum ($= 1 + \delta$) allowed by the sum rule. However, above the Fermi level, $n_c(\mathbf{k})$ is large ($= 1 - \delta$) all the way to the top of the band, whereas, it is zero for the other states. This costs considerable amount of kinetic energy. In the Schwinger boson case, the top of the band is empty, which is fine, but the kinetic energy is still large compared with E_{bound} because the Fermi energy is much higher, and because $n_c(\mathbf{k}) = 1 - \delta$ in the occupied region, smaller than value allowed by the sum rule, all the way to the bottom of the band.

c) *Hartree-Fock States in the Hubbard Model*: How are these results affected by the constraints? That there is an error due to the average treatment of the constraints can be seen from the average electron density

$$n_c = \sum_{\sigma} \langle c_{i\sigma}^{\dagger} c_{i\sigma} \rangle = \frac{1}{N} \sum_{\mathbf{k}} n_c(\mathbf{k}),$$

which does not equal its actual value $1 - \delta$ in the MF theory. Instead, it equals $(1 - \delta)^2$ in the Schwinger boson state, and $(1 - \delta^2)$ in the slave boson state. To assess the role of the constraints we consider the HF states of the Hubbard model in which constraints are treated exactly. We consider $U = \infty$ limit since mapping onto the $t - J$ model is exact (with $J = 0$). The HF ground state is ferromagnetic, with the up-spin band filled up to the holon Fermi level. The down-spin band is empty. This is the same ferromagnetic state found in SWB theory, and has the same fermi surface. However, now $n_c(\mathbf{k}) = n_{\uparrow}(\mathbf{k}) + n_{\downarrow}(\mathbf{k}) = 1$ in the occupied region, instead of $1 - \delta$. Since, as shown rigorously in section V., the states are essentially the same, the difference between the two results is a measure of the error (due to the constraints) in the Schwinger-boson theory.

Since the Fermi levels for the HF and Schwinger boson states coincide, the energies are proportional:

$$E_{sch} = (1 - \delta)E_{hub}.$$

Now, for all the states, $E_{kin} < 0$ at finite δ . Hence, E_{hub} is less than E_{sch} (and also less than E_{slave}) for all δ (see Fig. 2). At small δ , E_{hub} scales with δ , therefore the corrections due to constraints are small, i.e., of order δ^2 . To put it another way, since $E_{sch}/E_{hub} = 1 - \delta$, the effect of the constraints in the physically important small δ region is not large.

We see that kinetic energies of the MF and HF states are much larger than E_{bound} , the lowest possible KE without double occupancy. In the small δ region, E_{bound}/E_{MF} is between 1.5 to 2, i.e., the difference is comparable to the energy itself (Fig 3). This does not mean that E_{bound} corresponds to the actual state of the system. The key point is that $n_c(\mathbf{k})$ considerably smaller (by an amount $\propto \delta$ – the scale of KE) than the allowed value near the bottom of the band for the SWB and HF states, and considerably larger near the top of the band for the slave boson states. In the actual ground state, $n_c(\mathbf{k})$ is sure to depend on $\epsilon(\mathbf{k})$, and is likely to be close to $1 + \delta$ near the bottom, and close to zero near the top of the band. This means that fluctuations (in the MF states) that transfer electrons from near the top of the band to near the bottom will be dominant.

V. CONSTRAINTS: FROM SCHWINGER BOSONS TO SPIN WAVES

In this section we solve the constraint problem in the $U = \infty$ limit. It is convenient to use the functional integral technique. The Bose fields can be expressed as: $b_{i\sigma} = r_\sigma e^{i\phi_{i\sigma}}$. The constraint condition is given by

$$r_{i\uparrow}^2 + r_{i\downarrow}^2 + h_i^* h_i = 1,$$

which does not depend on the phases. The action is invariant under the gauge transformation: $b_{i\sigma} \rightarrow b_{i\sigma} e^{i\chi_i}$, $h_i \rightarrow h_i e^{i\chi_i}$. We can choose $\chi_i = -\phi_{i\uparrow}$, (or, equivalently select a gauge: $\phi_{i\uparrow} = 0$). The action is now independent of $\phi_{i\uparrow}$, which can be integrated out. The constraints are not effected. Next, we introduce the constraints into the functional integrals as δ functions, and use the latter to integrate out $r_{i\uparrow}$. This is equivalent to the replacement $b_{i\uparrow} = r_{i\uparrow} = (1 - r_{i\downarrow}^2 - h_i^* h_i)^{1/2}$ in the action. The remaining degrees of freedom define the generalized HP representation. It is easy to see that the corresponding Hamiltonian is obtained from the Schwinger boson Hamiltonian by using the following substitutions:

$$\begin{aligned} b_{i\downarrow} &\rightarrow b_i, & h_i &\rightarrow h_i, \\ b_{i,\uparrow}^\dagger &= b_{i,\uparrow} \rightarrow (1 - b_i^\dagger b_i - h_i^\dagger h_i)^{\frac{1}{2}}. \end{aligned}$$

The spin operators are represented as

$$\begin{aligned} S_i^+ &= (1 - b_i^\dagger b_i - h_i^\dagger h_i)^{\frac{1}{2}} b_i, \\ S_i^z &= \frac{1}{2} (1 - h_i^\dagger h_i) - b_i^\dagger b_i. \end{aligned}$$

In the absence of the holes, one recovers the usual HP representation for the Heisenberg ferromagnet. The generalization to the antiferromagnetic case is straightforward. As in the usual HP representation, the square-root represents an anholonomic constraint. This is of course not as stringent as the constraints in the Schwinger-boson (or slave-boson) representation. The electron operators are given by

$$\begin{aligned} c_{i\downarrow}^\dagger &= b_i^\dagger h_i \\ c_{i\uparrow}^\dagger &= (1 - b_i^\dagger b_i - h_i^\dagger h_i)^{\frac{1}{2}} h_i. \end{aligned}$$

In the last expression, we can drop the hole density operator within the square-root since it will not contribute (because h is fermionic). This gives, $c_{i\uparrow}^\dagger = (1 - b_i^\dagger b_i)^{\frac{1}{2}} h_i$. Since the constraints have been integrated out, there

is no gauge freedom left. Note that the b and h are gauge invariant since they are proportional to gauge-invariant operators S^+ and $c^\dagger$, respectively. As in the Heisenberg case, $b^\dagger$ creates a spin-wave excitation – a magnon. Similarly, the operator $h^\dagger$ creates a spinless hole excitation (not the unobservable holon of the SWB representation). Other difficulties associated with the constraints also disappear. For example, the local density of electrons is given by

$$\sum_{i\sigma} c_{i\sigma}^\dagger c_{i\sigma} = 1 - h_i^\dagger h_i,$$

which correctly yields, $n_c = 1 - \delta$, in contrast to, $n_c = (1 - \delta)^2$ in the Schwinger boson MF theory.

For $J = 0$ ($U = \infty$) the t - J Hamiltonian, in the generalized HP representation becomes

$$H = t \sum_{ij} [\{(1 - b_i^\dagger b_i)^{\frac{1}{2}}(1 - b_j^\dagger b_j)^{\frac{1}{2}} + b_i^\dagger b_j\} h_j^\dagger h_i + h.c.]. \quad (22)$$

Note that the total no of holes (N_h) and spin-wave excitations (N_b) are separately conserved. This follows from the conservation of the total electron number, $N_c = N - N_h$, and total magnetization $M_z = \sum_i S_i^z = \frac{1}{2}(N - N_h) - N_b$ in the parent Hubbard model. We will keep N_c fixed, which fixes N_h . But N_b is not fixed in the ordered state, but varies from 0 to N_c , where $N_b = N_c$ corresponds to zero magnetization.

We can diagonalize the Hamiltonian separately in each N_b subspace. The $N_b = 0$ subspace only has holes, and is of particular interest. In this case, $h_i = c_{i\uparrow}^\dagger$. All electrons have spin up, the average magnetization, $m = (1 - \delta)/2$, has the largest possible value. Within this subspace, the Hamiltonian reduces to

$$H_h = t \sum_{ij} (h_i^\dagger h_j + h.c.), \quad (23)$$

with no residual interaction. Note that this is nothing but the $U = \infty$ Hubbard Hamiltonian in the Hartree-Fock approximation in the fully spin-polarized subspace. In particular, it contains the ferromagnetic HF ground state (the generalized Nagaoka state) considered previously. We have thus shown that HF theory can be obtained from the Schwinger boson theory, and it takes the constraints into account exactly.

All wave functions corresponding to H_h are one-component Slater determinants, describing the charge sector. These are also exact wave functions of the full Hamiltonian (actually for all U) since in the $N_b = 0$ subspace, H_h is the exact Hamiltonian. It is not known rigorously whether the Nagaoka state is the true ground state at finite δ . A possible answer is that the true ground state belongs to a different N_b subspace, with magnetization $m = \frac{1}{2}(1 - \delta) - n_b$, which decreases continuously to zero δ_{cr} . Here, $n_b = N_b/N$ is the magnon density.

For finite n_b , we need to include the magnons. For small n_b , we expand the square root to leading order in n_b , which gives

$$H = t \sum_{ij} (h_i^\dagger h_j + h.c.) + t \sum_{ij} [(b_j^\dagger b_i - \frac{1}{2} b_i^\dagger b_i - \frac{1}{2} b_j^\dagger b_j) h_i^\dagger h_j + h.c.]. \quad (24)$$

The second term includes spin-wave effects, and extends the treatment of the $U = \infty$ Hubbard model beyond the Hartree-Fock level.

Note that there is no free magnon terms in the Hamiltonian. This is very different from the Heisenberg ferromagnet, where the lowest order ($1/S$) term is quadratic in bosonic operators. Here, magnons (spin) interact with holes (charge) at the lowest level. The simplest approximation is to do a MF decomposition of the quartic term, leading to quadratic Hamiltonians for holes and magnons: $H = H_b + H_h$. The hole Hamiltonian has the HF form (Eq. 23), but with a renormalized hopping parameter

$$t_h = t(1 + B - n_b),$$

where, $B = \langle b_i^\dagger b_j \rangle$. The bosonic mean-field Hamiltonian is given by

$$H_b = tD \sum_{ij} [b_i^\dagger b_j + b_j^\dagger b_i - b_i^\dagger b_i - b_j^\dagger b_j], \quad (25)$$

where $D = \langle h_i^\dagger h_j \rangle$. The magnon Hamiltonian has the same form as the one in the Heisenberg model. The magnon energy is then given by,

$$\omega_m(\mathbf{k}) = 2tD(\cos k_x + \cos k_y - 2). \quad (26)$$

It is easily shown that $D \leq 0$ is negative, so that $\omega_m(\mathbf{k}) \geq 0$. Therefore, there are no magnons at $T = 0$, and the Nagaoka state found in the HF approximation continues to be the ground state at this level of approximation. At finite T , magnons will proliferate just as in Heisenberg ferromagnet, and destroy long-range order. This is the correct result (Marmin-Wagner theorem). In principle, Takahashi's modified spin-wave method [15] can be used to describe the system at finite T .

Electron Green's Function and Missing Spectral Weight:

To obtain the Green's function for the up electrons, we expand the square-root. Now, at the MF level,

$$T_\tau \langle c_{i\uparrow}(\tau) c_{j\uparrow}^\dagger(\tau') \rangle = T_\tau \langle (1 - \frac{1}{2} b_i^\dagger(\tau) b_i(\tau) - \frac{1}{2} b_j^\dagger(\tau') b_j(\tau')) \rangle \langle h_i^\dagger(\tau) h_j(\tau') \rangle.$$

The average of the density operators simply gives the magnon density n_b , which is uniform. Hence, the Green's function is proportional to the hole Green's function:

$$G_{c\uparrow}(\mathbf{k}, \omega) = \frac{1 - n_b}{i\omega + \epsilon_h(\mathbf{k})}, \quad (27)$$

where $\epsilon_h(\mathbf{k}) = 2t_h(\cos k_x + \cos k_y) - \mu_h$ is the hole spectrum, which includes the chemical potential μ_h . This term is analogous to the condensate part in the Schwinger boson MF theory [16].

The Green's function for down electrons is a convolution of magnon and hole Green's functions, and is given by [17]

$$G_{c\downarrow}(\mathbf{k}, \omega) = \frac{1}{N} \sum_{\mathbf{q}} \frac{f(\epsilon_h(\mathbf{q})) + n(\omega_m(\mathbf{k} + \mathbf{q}))}{i\omega - \omega_m(\mathbf{k} + \mathbf{q}) + \epsilon_h(\mathbf{q})}. \quad (28)$$

This term is the analog of the non-condensate part in the SWB theory. The corresponding spectral function is nonzero only for $\omega > 0$ and is incoherent. It is absent in the HF theory. In the HF case there are two bands, described by simple poles. The up-spin band is the lower Hubbard band. The down-spin band is the upper Hubbard band which represents the doubly occupied sector and is separated by $U \rightarrow \infty$; it is absent in our case. In the simple HF theory, the spectral weights for each band is unity. This does not agree with the sum rule, which gives $1 + \delta$ and $1 - \delta$, respectively. The reason for the discrepancy is that magnons are missing in the simple HF theory. The expression for $G_{\downarrow}$ (Eq. 28), is precisely this (low-frequency) magnon contribution, with a total spectral weight of $n_b + \delta$. Together with the spectral weight of $G_{\uparrow}$, this gives a total spectral weight of $1 + \delta$, in agreement with the sum rule.

The Green's function in the spin-wave theory is very similar to that in the Schwinger boson theory [16]. However, in the latter case, magnetic ordering is along the x direction, so that the Green's function for up and down electrons are identical. For a proper comparison we need to take the average: $G_c = (G_{c\uparrow} + G_{c\downarrow})/2$. Although the structure of G_c in the two cases is the same, the spectral weight distribution between the coherent (pole) and incoherent pieces is different. For a comparison, it is convenient to express these in terms of the magnetization m which is the same in both theories, and equals $(1 - \delta)$ at $T = 0$. Recall that $n_c(\mathbf{k})$, and hence, the kinetic energy, is determined by the negative frequency part, i.e., by the pole piece ($G_{c\uparrow}$), which carries a spectral weight $= \frac{1}{2}(1 + \delta) + m$, which is greater than the (total) weight in the SB theory $= 2m$. In particular, it reduces to the HF value ($= 1$) at $T = 0$. Therefore, HF theory gives the best kinetic energy.

VI. CONCLUSIONS

In this paper we have shown that mean-field states of the t - J model exhibit complete spin-charge separation in the sense of Eq. (2), and in $d = 2$, have high kinetic energies. This supports qualitative arguments,

given in previous papers, that spin-charge separation localizes electrons which costs kinetic energy. The determination of E_{bound} is a key result since it includes excess KE which arises simply from the removal of double occupancy, but not necessarily from spin-charge separation. The calculation of $n_c(\mathbf{k})$ is also interesting, because it shows that mean-field values differ substantially from what is allowed by the sum rule at the band edges. Therefore fluctuations that transfer electrons from the occupied to the unoccupied regions, can, in principle, lower the energy significantly, i.e, by an amount comparable to KE itself.

The sum rule and the expression for E_{bound} are valid for arbitrary J . The kinetic energies presented here are for $U = \infty$ ($J = 0$) for which KE for MF and HF theories can be compared directly. Extension to finite J is not difficult since the MF problem has been solved exactly, but numerically. At finite J , the KE of the t - J model would be larger since the two terms compete. But conclusions of this paper are not going to change. Of course, J will be important to describe the spin singlets and for superconductivity.

The second important result is the mapping of the t - J model onto a generalized Holstein-Primakoff model. This allowed us to solve the difficult problem of the constraints for the SWB MF state at $U = \infty$. Our results show that constraints do not change the nature of the state – it remains fully spin-charge separated and has the same magnetic ordering. Extending the method to finite J is somewhat more complicated, but can be done. It will be useful to study fluctuations such as spin-charge recombination within the generalized HP model so as to take into account the constraints at the outset.

The mapping also establishes the connection between the SWB theory and the HF approximation. The latter has long been used to study magnetic ordering in strongly correlated systems. However, HF approximation fails at finite T (or, at finite frequency) because spin-waves are not included. For example, T_c at which magnetic order vanishes scales as U . The inclusion of the spin-waves via the mapping solves this problem, as spin-waves destroy LRO in 2-d at any finite T , and in $d = 3$ at a temperature $\sim t$ (as $U \rightarrow \infty$). It also demonstrates complete spin-charge separation in the HF plus spin-wave theory. This method will be useful in studying itinerant ferromagnetism in $d = 3$.

Our results suggest that fluctuations that result in spin-charge recombination are likely to be strong. Such processes lower KE by moving electrons from above the Fermi level to the bottom of the band. In previous papers we have treated recombination by RPA in which the physical electron appears as a fermionic collective mode [6]. In $d = 2$, it helps destroy the magnetic LRO, but, for δ not too large, the Fermi liquid state is not recovered. For finite J , additional spin-charge recombination between a pair of holons with preformed pair of spinons leads to d -wave superconductivity [7].

For the $U = \infty$ case, the destabilization of the HF state has been demonstrated by Shastry, Krishnamurthy and Anderson [18]. They considered the destruction of ferromagnetic LRO away from half-filling. Consider the ferromagnetic ground state with all spin up. Since $n_{k\uparrow} = 1$ in the occupied region, the only way kinetic energy can be lowered is by destroying an up spin electron at the Fermi level, and creating a down-spin electron at the bottom of the band. But creating a bare down-spin electron would cost U . Instead one must therefore create a projected down-spin electron. They showed that the MF state becomes unstable with respect to such spin modes. Note that in HP representation, (or SWB representation), the projected electron is a collective object created by $c_{\mathbf{k}\sigma}^\dagger = N^{-1/2} \sum_{\mathbf{q}} b_{\mathbf{k}+\mathbf{q},\sigma}^\dagger h_{\mathbf{q}}$. This process therefore corresponds to spin-charge binding, and is the same one considered by us.

References

- [1] P. W. Anderson, *Science* **235**, 1196 (1987).
- [2] P. W. Anderson, *Theory of Superconductivity in the High T_c Cuprates*, Princeton University Press, 1997.
- [3] C. Baskaran, Z. Zou and P. W. Anderson, *Solid. State. Commun.* **63**, 973 (1987).
- [4] M. Grilli and G. Kotliar, *Phys. Rev. Lett.* **64**, 1370 (1990); N. Nagaosa and P. A. Lee, *Phys. Rev. Lett.* **64**, 2450 (1990).

- [5] C. Jayaprakash, H. R. Krishnamurthy and S. K. Sarker, *Phys. Rev. B* **40**, 2610 (1989); D. Yoshika, J. Phys. Soc. Jpn. **58**, 3733 (1989).
- [6] S. K. Sarker, *Phys. Rev. B* **46**, 8617 (1992).
- [7] S. K. Sarker, *Phys. Rev. B* **61**, 8663 (2000).
- [8] S. Elitzur, *Phys. Rev. D* **12**, 3978 (1975).
- [9] S. K. Sarker, C. Jayaprakash, H. R. Krishnamurthy and W. Wenzel, *Phys. Rev. B* **43**, 8775 (1991); F. M. Hu, S. K. Sarker and C. Jayaprakash, *Phys. Rev. B* **50**, 17901 (1994).
- [10] D. P. Arovas and A. Auerbach, *Phys. Rev. B* **38**, 316 (1988).
- [11] S. K. Sarker, C. Jayaprakash, H. R. Krishnamurthy and M. Ma, *Phys. Rev. B* **40**, 5978 (1988); D. Yoshika, J. Phys. Soc. Jpn. **58**, 1516 (1989).
- [12] E. H. Lieb and F. Y. Wu, *Phys. Rev. Lett.* **20**, 1447 (1968).
- [13] M. Ogata and H. Shiba, *Phys. Rev. B* **41**, 2326 (1990).
- [14] This sum rule is known, and has been previously used in ref. [6]. However, the author is not aware whether a formal derivation has appeared in the literature.
- [15] M. Takahashi, *Phys. Rev. Lett.* **58**, 168 (1987); *Phys. Rev. B* **36**, 3791 (1987).
- [16] S. K. Sarker, C. Jayaprakash and H. R. Krishnamurthy, *Physica C* **228**, 309 (1994).
- [17] The expression for $G_{\downarrow}$ agrees with the spin-wave theory results derived by other methods, as it should. See, Yu. Irkhin and M. I. Katsnelson, *Sov. Phys.: Solid State Phys.* **25**, 1947 (1983); *J. Phys. C* **18**, 4173 (1985).
- [18] B. S. Shastry, H. R. Krishnamurthy and P. W. Anderson, *Phys. Rev. B* **41**, 2375 (1990).

FIGURE CAPTIONS

Fig.1:

Momentum distribution for various states as a function of energy. The slave boson case has a high kinetic energy because n_c is large all the way to the top of the band. For the other three states the top of the band is empty, but for the Schwinger boson case the energy is again large because n_c is small at the bottom of the band.

Fig.2:

Kinetic energy as a function of hole density δ . The HF energy is lower than either Schwinger boson or slave-boson case. However, E_{bound} is much lower than E_{bound} . The discrepancy between the HF and Schwinger boson energies measures the error due to poor treatment of constraints.

Fig.3:

E_{bound}/E_{hub} as a function of hole density. At small δ , the HF energy is larger by about a factor of two. (Note both energies are negative).





